

Intern.Journ. Pure and Appl. Math., 55, N4, (2009). 13-16.

Boundary integral equation for electromagnetic wave scattering by a homogeneous body of arbitrary shape

A G Ramm

Department of Mathematics
Kansas State University, Manhattan, KS 66506-2602, USA
ramm@math.ksu.edu

Abstract

Boundary integral equation is derived for the problem of scattering of electromagnetic waves by 3D homogeneous body of arbitrary shape.

MSC:45B05, 78A40, 78A45

PACS: 03.50.De

Key words: electromagnetic waves; wave scattering.

1 Introduction

Let $D \subset \mathbb{R}^3$ be a bounded domain with a smooth connected boundary S . The exterior domain $D' = \mathbb{R}^3 \setminus D$ is filled with a homogeneous material with parameters $(\epsilon_0, \sigma = 0, \mu_0)$, D is filled with a material with parameters $(\epsilon, \sigma, \mu_0)$, $\epsilon, \epsilon_0, \sigma, \mu_0$ are constants. Let $\epsilon' = \begin{cases} \epsilon + \frac{i\sigma}{\omega}, & \text{in } D, \\ \epsilon_0, & \text{in } D'. \end{cases}$ Here ω is the frequency, μ_0 is magnetic parameter, ϵ, ϵ_0 are dielectric parameters, $\sigma \geq 0$ is conductivity.

The governing equations in $\mathbb{R}^3$ are

$$\nabla \times E = i\omega\mu_0 H, \quad \nabla \times H = -i\omega\epsilon' E. \quad (1)$$

The boundary conditions on S are continuity of the tangential components of E and normal components of $\epsilon' E$ across S :

$$[N, E^+] = [N, E^-], \quad (2)$$

and

$$N \cdot (\epsilon' E^+) = N \cdot (\epsilon_0 E^-). \quad (3)$$

Here N is the unit normal to S , pointing into D' , $[N, E]$ ($E \cdot N$) is the cross (dot) product of vectors, and $E^+(E^-)$ is the limiting value of E as $x \rightarrow s \in S$, $x \in D(D')$.

We assume that the incident field (E_0, H_0) satisfies equations (1) in $\mathbb{R}^3$ with ϵ' replaced by ϵ_0 . For example one may take the incident field to be a plane wave: $E_0 = e_1 e^{ikx_3}$, where $e_i \cdot e_j = \delta_{ij}$, $\{e_j\}_{j=1}^3$ is the standard Euclidean basis, $\delta_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}$

If E is found then

$$H = (i\omega\mu_0)^{-1} \nabla \times E. \quad (4)$$

From (1) one gets

$$\nabla \times \nabla \times E - \mathcal{K}^2 E = 0 \quad \text{in } \mathbb{R}^3, \quad (5)$$

where

$$\mathcal{K}^2 = \begin{cases} K^2 & \text{in } D, \\ k^2 & \text{in } D', \end{cases} \quad \begin{aligned} K^2 &= \omega^2 \epsilon' \mu_0, \\ k^2 &= \omega^2 \epsilon_0 \mu_0. \end{aligned} \quad (6)$$

The scattering problem consists of finding the solution of equations (5), (2), (3), such that

$$E = E_0 + V, \quad (7)$$

$$V_r - ikV = o\left(\frac{1}{r}\right), \quad r := |x| \rightarrow \infty. \quad (8)$$

Assumption (8) means that V satisfies the radiation condition.

2 Derivation of the boundary integral equations

Let us look for the solution to (5), (2), (3), (7), (8), of the form:

$$E = \begin{cases} \nabla \times \int_S G(x, t) J(t) dt, & x \in D \\ \nabla \times \int_S g(x, t) j(t) dt + E_0(x), & x \in D', \end{cases} \quad (9)$$

where

$$g(x, y) = \frac{e^{ik|x-y|}}{4\pi|x-y|}, \quad G(x, y) = \frac{e^{iK|x-y|}}{4\pi|x-y|}. \quad (10)$$

The currents j and J are vector fields tangential to S .

Thus, there are four scalar unknowns: the two unknown vector fields j and J , tangential to S .

For any j and J vector E solves equation (5) in D and in D' and satisfies conditions (7) and (8) because g satisfies the radiation condition (8). Thus, (9) is the solution to the scattering problem if j and J can be chosen so that the boundary conditions (2), (3) are satisfied.

Condition (2) is a vector equation, which is equivalent to three scalar equations, and equation (3) is a scalar equation. Therefore equations (2) and (3) together are equivalent to four scalar equations for four unknown scalar functions, the coordinates of the tangential to S vector fields j and J .

Equation (2) can be written as a Fredholm-type integral equation. We have

$$\int_S [N_x, [\nabla_x G(x, t), J]]|_{x \rightarrow s, x \in D} = -\frac{A^+ J + J}{2} + \int_S \nabla_s G(s, t) N_s \cdot J(t) dt, \quad (11)$$

where the known formula for the limiting value of the normal derivative of the single-layer potential was used (see, e.g., [2]).

Since $J(s) \cdot N_s = 0$ (because $J(s)$ is a tangential vector field) and $J(s)$ is assumed Lipschitz, the last integral in (11) converges absolutely. Since we assume the surface sufficiently smooth, e.g., $S \in C^{1,a}$ $a > 0$, and the incident field is smooth, the currents j and J are as smooth as the data, in particular, they are Lipschitz. The class of surfaces, satisfying the condition $S \in C^{1,a}$, consists of surfaces whose graph in local coordinates is differentiable and its derivative satisfies the Hölder condition with the exponent $a \in (0, 1]$.

The operator A^+ in (11) is defined as

$$A^+ J = \int_S \frac{\partial G(s, t)}{\partial N_s} J(t) dt. \quad (12)$$

Thus, equation (2) can be written as:

$$\begin{aligned} -\frac{A^+ J + J}{2} + \int_S \nabla_s G(s, t) N_s \cdot J(t) dt &= -\frac{A^- j + j}{2} \\ &+ \int_S \nabla_s g(s, t) N_s \cdot j(t) dt + [N, E_0], \end{aligned} \quad (13)$$

where

$$A^- j = \int_S \frac{\partial g(s, t)}{\partial N_s} j(t) dt. \quad (14)$$

Equation (13) is of Fredholm type: the integral operators in (13) are compact in $C(S)$ and in $L^2(S)$. Equation (3) yields:

$$N_s \cdot \int_S [\nabla_x G(x, t), J(t)]|_{x \rightarrow s, x \in D} dt = N_s \cdot \int_S [\nabla_x g(x, t), j(t)]|_{x \rightarrow s, x \in D} dt + N_s \cdot E_0. \quad (15)$$

Equation (15) is singular.

Claim: *The integrals in (15) exist as Cauchy principal values.*

Let us verify this claim. One has

$$\begin{aligned} N_s \cdot [\nabla_x G(x, t), J(t)]|_{x \rightarrow D, x \in D} &= N_s \cdot \left[\frac{e^{iK r_{st}}}{4\pi r_{st}} \left(iK - \frac{1}{r_{st}} \right) r_{ts}^0, J(s) \right] \\ &\quad + O\left(\frac{1}{r_{st}}\right), \quad r_{ts}^0 := \frac{\vec{r}_{ts}}{r_{st}}, \quad r_{st} = |\vec{r}_{ts}|, \end{aligned} \quad (16)$$

because $|J(s) - J(t)| \leq c|s - t|$.

The singular term in (16) is

$$\frac{1}{4\pi r_{st}^2} N_s \cdot [r_{st}^0, J(s)] = \frac{|J(s)|}{4\pi r_{st}^2} \sin \theta, \quad (17)$$

where $\theta = \theta(s, t)$ is the angle between the x -axis and the vector $\vec{r}_{st}$. We choose the x -axis in the plane tangential to S at the point s so that it is directed along the vector $J(s)$.

Since

$$\int_0^\pi \sin \theta d\phi = 0, \quad t = e^{i\phi}, \quad 0 \leq \phi < 2\pi,$$

the **Claim** follows from Theorem 1.1 on p. 221 in [1]. This theorem says that a singular integral $\int_{\mathbb{R}^m} \frac{f(x, \theta)}{|x - y|^m} u(y) dy$ exists as a Cauchy principal value if

$$\int_{S^{m-1}} f(x, \theta) ds = 0, \quad \theta = \frac{y - x}{|y - x|}.$$

Numerical methods for solving Fredholm equations and singular integral equations are well developed ([1]). They are not discussed here.

References

- [1] S. Mikhlin, S. Prössdorf, Singular integra; operators, Springer-Verlag, Berlin, 1986.
- [2] C. Müller, Foundations of the mathematical theory of electromagnetic waves, Springer-Verlag, Berlin, 1969.